



CHESAPEAKE ENVIRONMENTAL PROTECTION ASSOCIATION, INC.
P.O. Box 117, Galesville, Maryland 20765

NEWSLETTER

Winter 2019-20

PRESIDENT'S MESSAGE

By Al Tucker



Are we funding our economy at the expense of ecosystem services from the environment?

In the very near future, climate change will radically alter the ecosystem and challenge its ability to provide the current level of service. Most of us want to feel financially secure. But, as with many financial decisions made on a daily basis, we make them

without thinking about their long-term costs. And with most of those decisions, we give even less thought to their impact on climate. There is emerging consensus among the public that climate change will significantly affect us, but little is understood how it will impact our economic well-being within a generation. At the national level, global warming results from the CO₂ emissions from fossil fuels, clearing of land by burning, or loss of forests by development. At the local level, the critical impacts will be driven primarily by land loss from development and sea-level rise. In each of these cases, a significant economic loss occurs from the loss of ecosystems services the land provided.

Our desire for modern technology drives the need for more energy. Unfortunately, the only immediately available form that can satisfy this demand comes from fossil fuels. Energy technologies that could supplant fossil fuels do not have the energetic quality or the economic and physical infrastructure to replace them. It appears that the demand for energy will outpace the ability of "clean energy" to stave off the effects of global warming if we continue with the business as usual approach. The economic value of land and forests that will be lost from sea-level rise and development will impose costs that will make maintaining the present level of quality-of-life difficult in future generations.

We will have to make tough choices that affect the energy delivery systems and the types of energy we consume. In 2018, the US saw the largest increase in energy consumption, in both absolute and percentage terms (~ 4%), since 2010. According to the International Energy Agency, the United States had the largest increase in oil and gas demand worldwide. The US consumption jumped 10% from the previous year. The annual increase in US demand last year was equivalent to the United Kingdom's current gas consumption.

Although it is publicly stated that the US consumed 11% of its energy from renewables, this number includes the burning of biomass. With the world-wide loss of forests to natural disasters, man-made destruction, and development, it is unlikely that the rate of biomass regeneration will ever come close to offsetting the increases in CO₂ production. The term renewable has a common perception that it includes only solar and wind, yet hydroelectric and geothermal contribute about 25% and hence only solar and wind contribute the remaining 30%, or about 4% of the total US consumption. It is highly unlikely that without a major intervention, solar and wind energy will ever have enough impact to reduce global warming to offset the anticipated sea-level rise.

The cost of the interventions and infrastructure required to change the pathway of global warming amounts to trillions of dollars. That is money that modern economies will find difficult to support unless the public demands change. To make these interventions, we will need to support national policies and be prepared to accept sacrifices.

Locally, we are paying for the loss of ecosystem services, caused by development. The stormwater fee pays for services that were previously provided by undeveloped land. Or consider the numerous failing septic fields in Anne Arundel County. (See article by Sally Hornor in this newsletter.) Individuals now bear the cost of compensating for the lost ecosystem services.

Sea-level rise will bring perhaps the largest loss of ecosystem services. Nuisance (or sunny day) flooding regularly plagues downtown Annapolis with the number of days expected to double in the next two decades. Businesses bear the costs of sandbagging and the increasing cost of their insurance premiums. Waterfront homeowners attempt to stave off sea-level rise with living shorelines or other means. These changes have hidden costs: decreased property values, reduced community investment, and lowered credit ratings, which leads to higher interest rates. Again, these costs represent direct payments of real money for services previously provided gratis by our environment.

In the near future, we will have to intervene and make hard choices. One unpalatable choice may be to limit the number of households in parts of the county and impose tighter, more costly regulations for environmental mitigation. We may need to impose stronger building restrictions where sea-level rise impacts the public infrastructure.

Keep in mind that economic well-being includes future financial security. The Council on Social Work Education defines economic well-being as:

"having present and future financial security. Present financial security includes the ability of individuals, families, and communities to consistently meet their basic needs ... Future financial security includes the ability to absorb financial shocks, meet financial goals, build financial assets, and maintain adequate income throughout the life-span."

IN THIS ISSUE

| | |
|--|----------|
| President's Message..... | 1 |
| Upgrading Septic Systems in AA County..... | 2 |
| Acidification of the Ocean and the Bay..... | 3 |
| Profile of a Trustee – Sally Hornor | 6 |

We can no longer defer to future generations the costs of borrowing from the ecosystem. A presentation at the 2016 CEPA forum by Elliot Campbell estimated that the value of ecosystem services for Anne Arundel to be approximately \$287,000,000 per year. Any land taken out of service will need to be replaced by man-made interventions that cost money. As more land is removed from ecosystem service, the remaining ecosystem service land becomes more costly to replace.

Not taking into account the economic contribution of ecosystem services provided to our economy at the national, state, and local levels needs to end immediately.

CEPA BOARD OF TRUSTEES

Albert Tucker, President
Bill Klepczynski, Vice-President
Lloyd Lewis, Secretary
Jeff Short, Treasurer

| | | |
|----------------|--------------------|------------------|
| Gary Antonides | Lee Greenbaum | Jerry Hill |
| Sally Horner | Anson (Tuck) Hines | Mike Lofton |
| Scott Knoche | Joan Turek | William Vosburgh |

UPGRADING SEPTIC SYSTEMS IN AA COUNTY

By Sally Horner

In Anne Arundel County and southern Md, there are two major sources of nutrients that flow into our rivers and into the Chesapeake Bay: stormwater and domestic wastewater. Anne Arundel County has done an admirable job in reducing the amount of nitrogen (N) that leaves wastewater treatment plants. They are also spending a lot of time and money trying to reduce the flow of stormwater, but our septic systems, or onsite sewage disposal systems (OSDS), are still a major source of nutrients, especially N. In fact, there are over 40,000 OSDS in the County and all of them leach N into surrounding soil. The closer the system is to the nearest waterway, either a tidal stream or river or the Bay, the more N it releases as there is less soil and plant growth to take up the N before it reaches the Bay.

To help meet its required nutrient reduction limits as set by the EPA Total Maximum Daily Loads (TMDL), AA County is trying to figure out how to reduce the input from OSDS. If a sufficient number of systems can be removed and the households hooked up to County sewer, the necessary reductions could be made. The problem with this plan is the cost: each system that is converted from septic to sewer will cost about \$60,000. Since about half of the OSDS in the County are in the Critical Area, being within 1000 feet of the water, conversion of just those 20,000 units would cost about \$1.5 billion.

This is the dilemma that was presented to the newly formed AA County Septic Task Force in 2017. Two CEPA trustees, myself and Lloyd Lewis, were asked to serve on this committee along with ten other County residents and a dozen County staff members. We met about 20 times over the following two years to see if we could come up with a solution that would allow the County to meet its TMDL and Watershed Implementation Plan (WIP) goals while still being affordable. While an earlier WIP (WIP II) required converting 20,000 OSDS, the 2019 WIP III called for only 5000 to 6000 systems to be converted, reducing the cost by 50%. This reduction in the number of required conversions was primarily due to the improved performance at County Wastewater Treatment Plants with respect to the reduction of nitrogen from sewage. Additionally, the earlier WIP called for the conversions to be done by 2025 while the new WIP

gives us until 2050. Other ways that the County can reduce nutrients include less expensive ways such as minor system upgrades and the use of Managed Aquifer Discharge (MAR). MAR was the key topic of our CEPA Forum in 2019 (see our website for individual presentations) and was described in the winter 2018-2019 CEPA newsletter which is also posted. Although an EPA official has recently stated that meeting the TMDL goals is not enforceable, County Executive Pittman has stated that we will continue to work toward these goals.

The 5000-6000 OSDS that are to be converted by 2050 include those that the County Health Dept. has determined to be public health risks. These systems are grouped into Onsite Wastewater Problem Management Areas; there are 33 in all and the top five areas that are given the highest priority for conversion are Clearview Village (north of Rt 100 in the Rock Creek watershed), Lake Placid area (Magothy River watershed), Amberley (Severn River watershed), and Edgewater Beach and Southdown Shores (South River watershed). Problem Area status is assigned based on a combination of several of the following factors: high water table, small lot size, impermeable soils, steep slopes, and setbacks from drinking water wells.

The County is hoping to convert about 200 septic systems to the sewer system per year in order to meet long term N reduction goals. The recommended process for such conversions will be a modification of the previous application process. As before, communities may petition for conversion and a simple majority vote (50% + 1) will trigger the requirement for all parcels in that community to be converted. Rather than charge the current front foot assessment to repay the County for its cost, a charge per household will be levied, thus eliminating the extra cost for larger properties or those with greater front footage.

According to AA County Bill 90-19, a payment deferment policy is recommended that would apply to households in the high-risk areas that meet at least one of the following criteria:

- parcels must be within the Critical Area,
- they must be in one of the Health Dept. Problem Areas as noted above,
- they must be in one of the areas slated for future sewer service as noted in the County Sewer and Water Master Plan or
- in an area adjacent to one of the areas listed above and also within the boundaries of a wastewater extension project that includes one of the criteria listed above.

If these criteria are met, residents may defer up to 50% of the total cost of connection and may have up to 40 years to pay it back. The ability to pay back in 40 years vs. 30 years was established by Bill 90-19 just passed by the County Council in December 2019. Deferred charges must be paid off in 40 years or within five years of the property being sold. A second bill proposed in December by Councilwoman Haire (95-19) would authorize the County to pay up to 25% of the total cost. At the Jan. 21 hearing of this bill, an amendment was passed that established that residents with a gross income of \$300,000 or less could receive up to 25% of the total cost while those making over \$300,000 could receive up to 12.5%. The County is willing to cover some of the cost of the conversions by these subsidies because the conversions will help them meet their TMDL goals.

Another topic discussed by the Septic Task Force is the need for public outreach and education to help residents understand the importance of converting septic systems to sewer. A key insight is the fact that households on septic systems contribute 18.6 pounds of N per year to our waterways whereas

households on sewer contribute 3 pounds N/year. Nitrogen is the primary nutrient that fuels algae blooms within our waterways, and algal blooms eventually lead to low dissolved oxygen concentrations in the Bay, causing loss of habitat for fish, crabs and oysters. This past December all of the major rivers in the county experienced a heavy bloom of dinoflagellates causing severe mahogany tides. Such events are triggered by nutrients flowing into our creeks and rivers from stormwater and septic systems.

One key concern expressed by Task Force members is the possible incentive to development that provision of sewer lines may permit. At the hearing for Bill 19-90, Councilwoman Haire estimated that about 100 new houses might be built in the County based on the extension of sewer lines. Since there are many criteria that must be met in order to qualify as a priority site for septic conversion, it is generally thought that the number of new houses will be limited.

There are still many steps that must be taken before this program is enacted, including the development of new policies relating to the timing of applications and coordination with State and County agencies. Stay tuned!

ACIDIFICATION OF THE OCEAN AND THE BAY

By Gary Antonides



In the Spring 2019 issue of their newsletter, *Catalyst*, the Union of Concerned Scientists reported that the surface waters of our oceans are nearly 30 percent more acidic than they were in 1850. If we don't reduce carbon emissions, they could be more than twice as acidic in 2100 than they were in 2000. Of the three major effects of increased carbon emissions, *ocean acidification* occurs the most quickly by far,

then *warming*, then *sea level rise*. If emissions decrease, ocean acidification will be the quickest to decrease. (<https://www.ucsusa.org/sites/default/files/attach/2019/05/Catalyst-Spring-2019.pdf>)

As carbon dioxide (CO₂) dissolves into the ocean, it forms carbonic acid, decreasing the ocean's pH. This is the process of ocean acidification, and it deprives shell-forming marine organisms (shellfish, corals, etc.) of the carbonate ions they need to build their protective shells. Acidifying waters eat away at the bottom of the world's food chain. When our global food chain is at risk, so are those who make their livings from, or subsist on, seafood.

The International Union for Conservation of Nature (IUCN), in <https://www.iucn.org/resources/issues-briefs/ocean-acidification>, says to combat acidification, CO₂ emissions need to be cut significantly and immediately and that conservation, restoration and permanent protection of at least 30% of the oceans are urgently needed.

The ocean absorbs over 25% of all human generated emissions from the atmosphere each year. As CO₂ dissolves in sea water and forms carbonic acid, it does so in parallel with ocean warming and deoxygenation. Interaction between these is often cumulative or even multiplicative, resulting in combined effects more severe than the sum of their individual impacts.

Present ocean acidity change is unprecedented in magnitude, occurring at a rate approximately ten times faster than anything

experienced during the last 300 million years. This rapid change is jeopardizing the ability of ocean systems to adapt to changes in CO₂ that naturally occur over millennia. Increases in ocean acidity will persist as long as concentrations of atmospheric CO₂ continue to rise. To avoid significant harm, IUCN says, atmospheric CO₂ levels need to get back to the 320-350 ppm range. A level of 400 ppm was exceeded in 2015.

Although more knowledge on the impacts of ocean acidification on marine life is needed, some predictions can be extrapolated from current knowledge. Some of the strongest evidence of the effects of ocean acidification comes from experiments on calcifying organisms. Increased sea water acidity has been shown to affect the formation and dissolution of calcium carbonate shells and skeletons in a number of marine species, including corals, oysters, mussels, and many phytoplankton and zooplankton species that form the base of marine food webs. Acidification can also cause changes in the growth and reproduction of marine species.

Increased ocean *temperatures* will also have direct effects on marine organisms and influence the geographical distribution of species. Some species such as reef-forming corals, are already living at their upper tolerance level, will have difficulty 'moving' fast enough to new areas. Changes in ocean temperature can also lead to coral bleaching, where corals expel the symbiotic algae living in their tissues, causing them to turn white.

The current emissions targets of the Intergovernmental Panel on Climate Change (IPCC) are to limit the global average temperature increase to less than 2°C. This is too much of a change to tackle the issue of ocean acidification. It will not reduce atmospheric CO₂ concentration to less than 400 ppm, and will significantly harm ocean life.

Another cause of ocean acidification is acid rain which contains nitrogen oxides and sulfur dioxide. They combine with atmospheric moisture to form acids. This is more concentrated near the coasts where the ocean is fed by rivers and runoff. (See "Acid Rain Has A Disproportionate Impact On Coastal Waters," Sept. 15, 2007, Woods Hole Oceanographic Institution <https://www.sciencedaily.com/releases/2007/09/070907175147.htm>). They say that the release of sulfur and nitrogen into the atmosphere by power plants and agriculture plays a minor role in making the ocean more acidic on a global scale, but the impact is greatly amplified in coastal waters. Coastal waters are already some of the most heavily affected parts of the ocean due to pollution, over-fishing, and climate change. Woods Hole found that the change in acidity in coastal waters due to nitrogen and sulfur was as much as 50 percent of the total change caused by CO₂. The most heavily affected areas tend to be downwind of power plants (particularly coal-fired plants) and, in the U.S., predominantly on the East Coast.

The EPA is working on all three causes of acidification: CO₂ emissions, acid-rain and excess nutrients. The part of the US Executive Branch called the Subcommittee on Ocean Science and Technology administers the [Interagency Working Group on Ocean Acidification](#), which includes 13 other agencies. The EPA collaborates with the National Estuary Program (NEP) which is deploying monitoring systems in estuaries around the country (San Francisco Bay, Santa Monica Bay, Tampa Bay, Massachusetts Bay, Casco Bay, Barnegat Bay, Long Island Sound, Corpus Christi Bay and Tillamook, OR). EPA is also measuring acidification in the Mid-Atlantic ocean offshore of Chesapeake and Delaware Bays and collaborating with states to study the interactions between nutrients and coastal acidification, and with universities to study impacts on shellfish.

An article in National Geographic, "Carbon dioxide in the water puts shelled animals at risk," April 27, 2017 (<https://www.nationalgeographic.com/environment/oceans/critical-issues-ocean-acidification/>) says that for tens of millions of years, Earth's oceans have maintained a relatively stable acidity level, but that this ancient balance is being undone by the recent and rapid drop in surface pH, which could have devastating global consequences. Scientists know that about half of the man-made CO₂ generated since the early 1800s has been absorbed over time by the oceans. This has reduced climate change from what it would have been if the CO₂ had remained in the air. However, relatively new research is finding that the massive amounts of CO₂ in the oceans is affecting the life cycles of many marine organisms, particularly those at the lower end of the food chain.

The pH scale runs from 0 to 14, where solutions with low numbers are acidic and those with higher numbers are basic. Seven is neutral. As examples, lye has a pH of 13 and lemon juice has a pH of 2. Over the past 300 million years, the average ocean pH has been slightly basic (about 8.2). Today, it is around 8.1. A drop of 0.1 might not seem like a lot, but the pH scale is logarithmic and that drop amounts to about 25%. (pH 4 is ten times more acidic than pH 5) If we continue to add CO₂ at current rates, seawater pH may drop to 7.8 or 7.9 by the end of the century (an increase in acidity of *another* 60 to 100 percent), creating an ocean more acidic than it has been for the past 20 million years or more.

A Smithsonian website, <https://ocean.si.edu/ocean-life/invertebrates/ocean-acidification> says it was once thought that ocean acidification might be a good thing because it leaves less CO₂ in the air. But in the past decade, we've realized that this has come at the cost of changing the ocean's chemistry. Also, scientists once assumed that rivers carried enough dissolved chemicals from rocks to the ocean to keep the ocean's pH stable (called "buffering.") But so much CO₂ is dissolving into the ocean that this natural buffering hasn't been able to keep up.

Scientists have been tracking ocean pH for more than 30 years, but biological studies really only started in 2003. Some organisms will survive or even thrive under more acidic conditions while others will struggle to adapt, or go extinct. Field studies are difficult because of other changes, like warming.

Living things can be very sensitive to small changes in pH. In humans, for example, normal blood pH is about 7.4. A drop in blood pH of 0.2-0.3 can cause seizures, comas, and even death. A small change in the pH of seawater can also impact communication, reproduction, and growth of marine life. The pH of the ocean fluctuates somewhat as a result of natural processes, and ocean organisms are able to survive these normal changes, but not the rapid recent changes. During the last great acidification event 55 million years ago, there were mass extinctions in some species including deep sea invertebrates. The rise in seawater acidity of 25 percent that we already have is already affecting some ocean organisms.

Oysters, Mussels, Urchins And Starfish - Generally, shelled animals, including mussels, clams, urchins and starfish, are going to have trouble building their shells in more acidic water. Mussels and oysters are expected to grow less shell by 25 percent and 10 percent respectively by the end of the century. Oyster larvae will fail to even begin growing their shells. Urchin and starfish shells dissolve even more quickly than corals.

A Yale report, "Northwest Oyster Die-offs Show Ocean Acidification Has Arrived," by Elizabeth Grossman, 11/21/11

(https://e360.yale.edu/features/northwest_oyster_die-offs_show_ocean_acidification_has_arrived) says that In Netarts Bay in Oregon, from 2006 to 2008, oyster larvae began dying, with losses of 70 to 80 percent. Burke Hales, from Oregon State University, determined that the cause was acidic seawater. Because of patterns of ocean circulation, colder, more acidic waters stream ashore in the fjords, bays, and estuaries of Oregon, Washington, and British Columbia. The corrosive waters prevented oyster larvae from forming shells. Oysters elsewhere in the Pacific Northwest have also experienced reproductive failure. Some of the largest operations are now buffering the water in which they grow their larvae, giving their tanks a dose of antacid in the form of sodium bicarbonate.

Problems with ocean acidification are also starting to be seen on the U.S.'s Atlantic coast. Agricultural runoff and sewage already take a toll on the once-thriving oyster business in the Chesapeake Bay, and now rising ocean acidity is further exacerbating the problems. However, studies have found that, for some reason, crustaceans such as lobsters, crabs, and shrimp grow even stronger shells under higher acidity.



Zooplankton - Many species of zooplankton (tiny drifting animals) build shells, such as these sea butterflies in the Arctic. They are small, but almost all larger life eats zooplankton or other animals that eat

zooplankton. These tiny organisms reproduce so quickly that they may be able to adapt to acidity better than slow-reproducing animals, but experiments have found that some of their shells do dissolve rapidly in acidic conditions.



Plants and Algae - Plants and many algae may thrive with acidic conditions. Seagrasses are nurseries for many larger fish, and can be home to thousands of different organisms. Unfortunately, they are in decline for a number of other reasons, especially pollution

flowing into coastal seawater. It's unlikely that a boost in acidification will compensate entirely for that.

One major group of phytoplankton (single celled algae) grows shells. Early studies found that their shells weakened with increased acidity. But a longer-term study let them reproduce for 700 generations (taking about 12 months) in the warmer and more acidic conditions expected to become reality in 100 years, and they were able to adapt, growing strong shells.



Fish and Jellyfish - While fish don't have shells, acidification changes the pH of the fish's blood (acidosis). A small change in pH can make a huge difference in survival by affecting its catching and digestion of food, its growth, its avoidance of predators, and its reproduction. Clownfish can normally hear and avoid

predators, but in more acidic water, they do not flee threatening noise. They also stray farther from home and have trouble getting back.

One big unknown is whether acidification will affect jellyfish. They compete with fish and other predators for food. If jellyfish thrive under warm and more acidic conditions while most other organisms suffer, it's possible that jellies will dominate in some places, a problem already seen in parts of the ocean.

Coral Reefs - Acidification corrodes existing coral skeletons as well as slows the growth of new ones. A recent study predicts that by roughly 2080 ocean conditions will be so acidic that even otherwise healthy coral reefs will be eroding more quickly than they can rebuild. Acidification also impacts coral animals, but at this point only a few coral species have been studied.



Studying Acidification - Geologists study the potential effects of acidification by digging into Earth's past when ocean CO₂ and temperature were similar to conditions found today. One way is to study cores, soil and rock samples taken from deep in the Earth's crust, with layers that go back 65 million years. The chemical composition of fossils in cores from the deep ocean show that it's been 35 million years since the Earth last experienced today's high levels of atmospheric CO₂. But to predict what might

happen by the end of the century, geologists have to look back another 20 million years. Some 55.8 million years ago, massive amounts of CO₂ were released into the atmosphere, and temperatures rose by about 9°F. We don't know why, but, like today, the pH of the deep ocean dropped quickly, and so much of the shelled sea life disappeared that the sediment changed from primarily white calcium carbonate "chalk" to red-brown mud. Today, CO₂ levels are rising even faster than then.

Another way to study acidification is to perform controlled laboratory experiments, but other stressors such as warming and pollution have to be considered. Short-term studies might not uncover the potential for some species to adapt to decreasing ocean pH. For example, in one test, a deepwater coral showed a significant decline in its ability to maintain its skeleton during the first week of exposure to decreased pH, but after six months, the coral had adjusted and returned to a normal growth rate.



This photo shows CO₂ bubbles from volcanic vents in the reef off the coast of Papua New Guinea. One effect of these CO₂ seeps was that large boulder corals replaced complex branching forms and, in some places, the corals disappeared entirely.

There are places scattered throughout the ocean where CO₂-rich water from vents like these lower the pH in surrounding waters. Scientists study these for clues as

to what an acidified ocean will look like. Researchers working off the Italian coast compared the ability of 79 species of bottom-dwelling invertebrates to settle in areas at different distances

from CO₂ vents. For most species, including worms, mollusks, and crustaceans, the closer to the vent, the fewer individuals were able to colonize or survive. Reef-building corals, snails, barnacles, sea urchins, and coralline algae were absent or much less abundant in the acidified water, which was dominated by dense stands of sea grass and brown algae.



This photo shows a test tube that is 60-feet deep and holds almost 15,000 gallons of water. Extra CO₂ can be added, making the water inside more acidic. Studies with these "mesocosms" can include many other

effects beyond acidification, such as warming, pollution, and overfishing. Scientists from five European countries built ten of these and placed them in a Swedish fjord. After letting plankton and other tiny organisms drift or swim in, they sealed the test tubes and decreased the pH to 7.8, the expected acidity for 2100, in half of them to see if and how the organisms adapt. They have not yet completed the experiment, but, if successful, it can be repeated in different ocean areas around the world.

Chesapeake Bay - "Growing acidification of the Chesapeake Bay threatens crabs, oysters, other life," by Scott Dance, Baltimore Sun, Oct. 5, 2017 reports that, during recent summers, researchers aboard a University of Delaware research vessel collected water samples from the mouth of the Susquehanna River to Solomons Island in order to find out when and where the waters of the Chesapeake Bay were turning most acidic. One finding was that acidification is likely to come faster to the Bay than to our oceans. Researchers are beginning to realize that acidification will compound the ecological challenges already affecting the bay.

Experiments are showing that blue crabs, marsh grasses and algae could theoretically thrive in the conditions expected to develop over the next century. But the acidification is a threat to other bay species, such as oysters, which are a key source of food for crabs. Scientists say acidification could dramatically alter the delicate balances in the bay ecosystem. With so many things affecting bay creatures (acidity, warming, pollution), research is complicated.

The University of Delaware research found a zone of increasing acidity at depths of about 30 to 50 feet across the Bay. While surface waters hover around the pH norm of 8.2, the deeper waters registered almost one point lower (10 times more acidic). The researchers believe it's not only the global effects of CO₂ emissions, but also the dead zones of low or no oxygen that are created when nitrogen and phosphorus runoff from farms and lawns, and fertilize large algae blooms. Microbes strip oxygen from the water when they die and decompose, and release more CO₂ in the process. When the water is already stripped of oxygen, and organic matter decomposes, the bacteria use up other compounds in the water which produce hydrogen sulfide, which makes the muck in the bay smell like rotten eggs.

Others are testing what could happen to the Bay's crabs and oysters. For her dissertation, UMCES doctoral candidate Hillary Glandon exposed blue crabs to both warmer and more acidic waters and found that acidification alone didn't affect them, but when coupled with warmer waters, crabs grew faster, and ate more food. While crabs could thrive in warmer and more acidic water, the oysters and mussels they eat could be struggling.

Looking to the Future - When CO₂ in the atmosphere passed 400 parts per million (ppm) in 2015, it became higher than at any time in the last one million years (maybe even 25 million years). The "safe" level is around 350 ppm, a milestone we passed in 1988. Without ocean absorption, atmospheric CO₂ would be closer to 475 ppm. The most realistic way to lower this number or to keep it from getting astronomically higher would be to reduce our carbon emissions by burning less fossil fuels and finding more carbon sinks, such as growing mangroves, seagrasses, and marshes. If we did this, it would still take over hundreds of thousands of years for CO₂ in the atmosphere and ocean to stabilize again. Even if we stopped emitting all carbon right now, ocean acidification would not end immediately because there would be a lag before we see the effects. The climate would continue to change, the atmosphere would continue to warm and the ocean would continue to acidify. The main effect of increasing CO₂ that weighs on people's minds is global warming. Some proposals suggest we reflect sunlight back into space from the atmosphere. However, this solution does not remove CO₂ from the atmosphere, or keep it from dissolving into the ocean.

**PROFILE OF A TRUSTEE
SALLY HORNOR**



Sally's love of coastal ecosystems was no doubt inspired by growing up on the Atlantic seaboard. She majored in Biology at Goucher College (Towson MD) and spent college summers at Woods Hole Oceanographic Institution in Massachusetts. She then went on to earn an MS in Microbiology and a Ph.D. in Ecology, both at the Univ. of Connecticut. She has since worked as an aquatic microbial ecologist in such diverse ecosystems as sewage sludge, bog lakes, rivers and the Chesapeake Bay watershed.

After a post-doc fellowship at SUNY Syracuse, Sally joined the faculty at Virginia Tech where she taught limnology and aquatic microbiology. Sally and her husband, Tom Caperna, moved to Arnold MD in 1984 when Tom accepted a research scientist position at USDA in Beltsville. While raising two children, Sally taught part-time at Anne Arundel Community College, and then taught full time in 1993, teaching primarily ecology and microbiology; retiring from teaching in 2015. While at AACC, she was the scientific director of Operation Clearwater, a program started about 50 years ago by the Severn River Association to monitor the bacterial water quality of bathing beaches and marinas during the swimming season. She expanded this program in partnership with the Magothy River Association, the South River Federation, the West and Rhode Riverkeeper and the Advocates for Herring Bay so that bacterial water quality monitoring of all of the major tributaries in AA County were included. She has also worked with the Severn River Association and the Magothy River Association to restore historic oyster bars and to encourage citizen science projects in water quality monitoring and monitoring of submerged aquatic vegetation. Sally is also on the board of the Alliance for the Chesapeake Bay and the Magothy River Land Trust as well as serving on the AA County Environmental Commission.

Sally and her husband get out on the Chesapeake Bay sailing and kayaking as much as possible in the summer and travel to visit a diversity of ecosystems as time permits. Their grown children and two grandchildren live in Baltimore and Virginia.

To join CEPA, please fill out the form below and send it with your check to:
CEPA, PO Box 117, Galesville, MD 20765
or join online at www.cepaonline.org.



CEPA MEMBERSHIP

A CEPA membership entitles you to receive our newsletter and to vote for our Trustees.

Name _____
 Address _____

 Phone _____
 Email _____

Enclosed is:

- \$30. for my CEPA membership
- \$50. for my Sponsoring CEPA membership
- \$100. for my Sustaining CEPA membership

Newsletters:

- Please send me emails when the newsletters are posted.
- Please send hard copies of the newsletters to me by mail



CEPA
P.O. Box 117
Galesville, MD 20765